

a 62445



WHO/Ma1/358
14 September 1962

ORIGINAL: ENGLISH

THE ANOPHELES GAMBIAE COMPLEX

by

G. Davidson
Ross Institute of Tropical Hygiene, London

The existence of two fresh-water forms (A and B) of Anopheles gambiae Giles which, when crossed, produce sterile F_1 males has been established by Davidson & Jackson.¹ A salt-water tolerant strain of the same species from near Tanga, Tanganyika, has been shown to be partially incompatible with two strains of the Group A fresh-water form,^{2,3} while the variety melas has been shown to be incompatible with a fresh-water form (group unknown) from a nearby area in Liberia.⁴

Now, the Tanga salt-water form and the A. melas from Liberia have been crossed with each other and with each of the two fresh-water forms. Table 1 enumerates these crosses and gives the proportion of male to female offspring in the F_1 generation.

All the crosses produced sterile F_1 males. Sterility was established in three ways:

- (1) by microscopic examination of the testes;
- (2) by allowing interbreeding of F_1 males and females in cages and observing the eggs laid by the females;
- (3) by artificially mating F_1 males and females after the method of Baker et al.⁵

The degree of atrophy of the testes varied considerably. In some, the testes were reduced to an organ barely distinguishable from the vas deferens and in which no sign of spermatogenesis could be distinguished. In others the testes appeared normal in size and showed all stages of spermatogenesis up to what appeared to be normal, tailed spermatozoa. These, however, showed no sign of movement in physiological saline. All F_1 adults appeared normal in size and

vigour and F_2 eggs were obtained from all but two of the crosses. These eggs were almost invariably sterile and microscopic examination showed no signs of embryonic development. Only on two occasions did a very few larvae hatch, and these were from the reciprocal crosses between the Pare fresh-water form and the Tanga salt-water form.

An excess of males was always produced from crosses involving fresh-water form males and salt-water form females (both the Tanga strain and A. melas). This was most marked in the fresh-water form male x melas female cross. The two crosses between fresh-water form females and Tanga salt-water form males produced a slight excess of females in the F_1 generation. Similar crosses between fresh-water females and melas males gave a slight excess of males. The reciprocal crosses between the Tanga salt-water form and A. melas resulted in near-normal sex-ratios.

That some at least of the hybrid females were reproductively normal was shown by the production of viable offspring from backcrosses. These offspring all showed near-normal sex-ratios (Table 2). In addition to those backcrosses given in this table, the backcross Tanga salt-water form male x hybrid Tanga male/Pare female was also successful, but no sex-ratio was recorded.

It would thus appear that Anopheles gambiae is a complex of at least four partially incompatible forms. Whether any or all of these deserve rank as separate species remains to be seen. If complete absence of gene-flow constitutes a criterion of specific rank, then these forms cannot be considered as separate species. A restricted flow is possible through backcrossing, at least in the artificial conditions of the laboratory.

REFERENCES

1. Davidson, G. & Jackson, C. E. (1962) Bull. Wld Hlth Org. (In press)
2. Annual Report of East African Institute of Malaria and Vector-Borne Diseases, 1960, East Africa High Commission, Nairobi, Kenya
3. Patterson, H.E. (1962) Nature, 195, 469
4. Annual Report of Liberian Institute of American Foundation for Tropical Medicine, 1960, Harbel, Liberia
5. Baker, R. H., French, W. L. & Kitzmiller, J. B. (1962) Mosquito News, 22, 16

TABLE 1. CROSSES BETWEEN FRESH-WATER AND SALT-WATER
FORMS OF ANOPHELES GAMBIAE

Parents			F ₁ Generation	
Male	Female	Total	Percentage male	Percentage female
Lagos (A)	TSW	66	61	39
Kisumu (A)	TSW	250	56	44
Pare (B)	TSW	156	56	44
Bobo (B)	TSW	138	60	40
TSW	Lagos (A)	1010	42	58
TSW	Pare (B)	998	36	64
Kisumu (A)	<u>melas</u>	113	76	24
Liberia (A)	<u>melas</u>	326	99	1
Pare (B)	<u>melas</u>	122	81	19
<u>melas</u>	Diggi (A)	94	51	49
<u>melas</u>	Pare (B)	88	60	40
<u>melas</u>	TSW	62	45	55
TSW	<u>melas</u>	193	57	43

- Lagos - Lagos, Nigeria
 Diggi - Diggi, W. Sokoto, N. Nigeria
 Kisumu - Kisumu, Kenya
 Pare - Pare, Tanganyika
 Bobo - BoboDioulasso, Upper Volta
 Liberia - Kpain, Liberia
 TSW - Tanganyika salt-water form
melas - Anopheles melas, Liberia
- A - group A) see Davidson
 B - group B) and Jackson¹

TABLE 2. BACKCROSSES BETWEEN FRESH-WATER AND
SALT-WATER FORMS OF ANOPHELES GAMBIAE¹

Parents			F ₁ Generation	
Male	Female Hybrid	Total	Percentage male	Percentage female
TSW	<u>melas</u> ♂ / TSW ♀	134	47	53
<u>melas</u>	<u>melas</u> ♂ / TSW ♀	20	45	55
Diggi	<u>melas</u> ♂ / Diggi ♀	305	46	54
<u>melas</u>	<u>melas</u> ♂ / Pare ♀	19	37	63
<u>melas</u>	Pare ♂ / <u>melas</u> ♀	21	57	43

¹ See footnotes of Table 1

The purpose of the WHO/Mal series of documents is threefold:

(a) to acquaint WHO staff, national institutes and individual research or public health workers with the changing trends of malaria research and the progress of malaria eradication by means of summaries of some relevant problems;

(b) to distribute to the groups mentioned above those field reports and other communications which are of particular interest but which would not normally be printed in any WHO publications;

(c) to make available to interested readers some papers which will eventually appear in print but which, on account of their immediate interest or importance, deserve to be known without undue delay.

The issue of a paper in this series does not therefore constitute formal publication and a paper so issued may, with the agreement of the author and WHO, be published in a WHO periodical or elsewhere.

Authors alone are responsible for views expressed in signed articles. The mention of manufacturing companies or of their proprietary products does not imply that they are recommended or endorsed by the World Health Organization.