




2 65159

A FIELD TRIAL USING DICHLORVOS IN  
 MAMASANI AREA, KAZEROUN, SOUTHERN IRAN. 1965-1966

by

B. Samimi, M. Motabar, R. Rouhani, and M. Mottaghi



The Editors regret that by an oversight, for which the authors are in no way responsible, the proportions of parous (NB - read "parous" for "porous") mosquitos, shown in Table 5, page 8 are incorrectly expressed. The following text should therefore be substituted:

TABLE 5. PROPORTION PAROUS

		<u>A. stephensi</u>		<u>A. superpictus</u>	
		Before application of dichlorvos	After application of dichlorvos	Before application of dichlorvos	After application of dichlorvos
1965	Control village	0.17	0.21	0.67	-
	Treated village	0.63	0.091	0.55	0.26
1966	Control village	-	0.44	-	0.28
	Treated village	-	0.074	-	0.28

The issue of this document does not constitute formal publication. It should not be reviewed, abstracted or quoted without the agreement of the World Health Organization. Authors alone are responsible for views expressed in signed articles.

Ce document ne constitue pas une publication. Il ne doit faire l'objet d'aucun compte rendu ou résumé ni d'aucune citation sans l'autorisation de l'Organisation Mondiale de la Santé. Les opinions exprimées dans les articles signés n'engagent que leurs auteurs.