

## INDEX

- Acidosis, metabolic, 49-50  
Activity, and food intake, 487  
Adaptation, metabolic, to nutrient supply, 451  
Adenoma, 86  
Affluence, nutritional problems of, 482-499  
Aflatoxin, and neoplasm, 184  
  in groundnut and cottonseed cake, 394  
Alcohol, as energy source, 458  
Alcoholism, and magnesium deficiency, 155  
  and pellagra, 127  
  and thiamine deficiency, 136, 477  
  in urban immigrants, 245  
Amino acids, biological value, 464  
  chemical scoring, 463  
  essential, serum level in PEM, 35  
  food fortification with, 381, 383, 385  
  imbalance, and pellagra, 126  
  limiting, definition of, 462-463  
  ratio of non-essential to essential, 569  
Aminoaciduria, in vitamin D deficiency, 115  
Anaemia, megaloblastic, 151  
  *see also* Folate deficiency; Vitamin B<sub>12</sub> deficiency  
Anaemia, nutritional, 55-79  
  definition, 55, 62  
  determination, methods, 69-71  
  diagnosis, 69-71  
  etiology, 60-61  
  feasibility of control, 18  
  prevalence, 18, 62-64  
  prevention, 73-74  
  social significance, 18  
  symptoms, 68  
  treatment, 71-73  
    iron supplementation, 315, 316  
  and infections, 203  
  and parasitism, 61-62  
  and PEM, 33  
  *see also* Iron deficiency; Folate deficiency  
Anaemia, pernicious, 67, 72, 478  
Anthropometry, 262, 414  
  indicator of nutritional status, 501, 536-546  
  methods, accuracy, 546  
  weight charts, 520-529  
  *see also* Skinfold thickness  
Antibody formation, and nutrient deficiency, 199-200  
  in anaemia, 68  
  in PEM, 271  
Anticonvulsants, and rickets, 112  
Antivitamins, deoxypyridoxine, 149, 199  
  omega-methylpantothenic acid, 199  
  thiaminase, 144  
Arm circumference, 545  
  *see also* Anthropometry  
Ascariasis, and PEM, 43  
Ascorbic acid, food fortification with, 383  
  losses during food storage, 357, 358  
  pharmacological effects, 479  
  recommended daily intake, 121, 456  
  supplements in times of disaster, 413  
  synthesis from glucose, 121  
  and hydroxyproline synthesis, 200  
Ascorbic acid deficiency, 479  
  and antibody formation, 199  
  *see also* Scurvy  
Atherosclerosis, 340, 489  
  relation to diet, 490-491  
  *see also* Cardiovascular disease  
Auxiliary health workers, in rural areas, 234  
  role as nutrition educators, 290-292  
  training and supervision, 265, 281  
Avitaminoses, *see* Vitamin deficiencies
- Baby foods, fortification, iron, 74  
  vitamin D, 115  
  local multimixes, 48  
  vitamin E content, 151  
  and folate deficiency, 74  
  and infantile scurvy, 123  
  *see also* Infant foods; Weaning foods  
Balances, for infants and toddlers, practical considerations, 525-526  
BCG-vaccine, *see* Tuberculosis  
Beans, iron availability, 58  
  vitamin B content, 476  
Beef, vitamin B content, 476  
Beriberi, 13  
  diagnosis, transketolase activity, 139  
  epidemiology, 140-144  
  mortality, morbidity, Japan, 141-143  
  symptoms, 136-145  
    adults, 137-139  
    infants, 139-140  
  treatment and prevention, 144-145  
  and infections, 198  
Bicarbonate, and acidosis in diarrhoea, 49  
  and rehydration solutions, 50-51

- Biometry (biometrics)**, 179-185  
 definition, 181  
 histopathological examination, 184  
 serological examination, 184
- Biotin deficiency, and antibody formation**, 199
- Birth rates, and maternal and child health**, 229  
 and population structure, 165
- Birth weight, distribution in different countries**, 508, 510  
 indicator of nutritional status, 501, 507-510  
 influence of social structure on, 509  
 low, definition, 509  
 and spacing of pregnancies, 229
- Bitot's spots**, 99, 100, 103-104
- Blindness, and vitamin A deficiency**, 96, 97
- Blood, calcium level in rickets**, 114-115  
 serum albumin in scurvy, 124
- Body weight, criterion for classification of PEM**, 532-533  
 recommendations for (USA), 455-456, 486
- Bone, and calcium metabolism**, 469-470
- Bottle feeding, dangers**, 30, 226  
 use in urban Africa, 248  
*see also* Infant feeding; Baby foods
- Bread, consumption in urban Africa**, 247  
 iodine fortification, 90  
 iron and vitamins fortification, 383  
 and dental caries, 303
- Breast feeding**, 46, 230  
 inadequate, 30, 308  
 recommendations concerning, 391  
 social aspects of, 226  
 and birth spacing, 48  
 and child malnutrition, 30  
 and working mothers, 304  
 in urban immigrants, 248, 390  
*see also* Lactation
- Breast milk, biological protein value**, 464  
 protein content, 462  
 supplements, *see* Weaning foods  
 thiamine content, 141, 477  
 vitamin A content, 474
- Browning, nonenzymatic**, 358
- Butter, vitamin A content**, 95
- Calcium**, 469-470  
 food fortification with, 383  
 interference with magnesium, 155  
 loss in tuberculosis, 202  
 metabolism and rickets, 112-118  
 recommended daily intake, 455, 470  
 relation to phosphorus, 470  
 requirement, adaptation to intake, 451, 469
- Capillaria philippinensis*, 77
- Carbohydrate metabolism, and obesity**, 484
- Cardiovascular disease**, 489-494  
 epidemiology, 489  
 etiology, 491  
 risk factors, 489-492  
 overnutrition, 341, 485  
 serum lipid level, 489  
 trace elements, 156  
 and thiamine deficiency, 137-138
- Caries**, *see* Dental caries
- $\beta$ -Carotene**, 95, 109, 472  
 conversion rate to vitamin A in PEM, 98, 473  
 food fortification with, 384
- Cassava, vitamin B content**, 476
- Cell proliferation, and growth**, 196
- Cereals, consumption in developing countries**, 15  
 consumption in North America, 15  
 expenditure on, in different regions, 362  
 iodine source, 84  
 new varieties, 345-346  
 price control, 366  
 storage, nutritional loss, 358  
 vitamin B content, 476  
 in weaning food mixtures, 393
- Cheese, vitamin B content**, 476
- Chemical score**, 462-465
- Chickenpox, and kwashiorkor**, 29
- Childhood mortality**, 183  
 age-specific ratios, 506  
 control, 215  
 data collection, urban areas, 251  
 factors responsible for, 16  
 diarrhoea, 203  
 infections, general, 209  
 measles, 207-208  
 PEM, 28, 44, 258, 322  
 vitamin A deficiency, 97  
 indicator of nutritional status, 502-507  
 and birth order, 229, 232  
 in urban immigrants, Africa, 249  
*see also* Mortality rates
- Children, age range in the family**, 225-227  
*see also* Preschool children
- Cholecalciferol**, *see* Vitamin D
- Cholesterol, dietary, and cardiovascular disease**, 489, 493

- Chromium, 156, 471
- Cobalt, 156, 471
- Cod liver oil, vitamin A supplement, 107  
vitamin D supplement, 474
- Celiac disease, 77  
and iron absorption, 60
- Collagen, 200
- Conjugase (gamma-glutamylpeptidase), 63
- Contamination, of foods, 357  
prevention, 359-360
- Contraceptives, oral, and vitamin B<sub>6</sub> deficiency, 150
- Copper, 156, 471-472
- Corn-soya-milk (CSM), 301, 399
- Cortisone, and infections, 201
- Coryza, 259-260
- Cost-effectiveness, food and nutrition policies, 435  
nutrition programmes, 219, 221  
nutrition rehabilitation centres, 329-330  
supplementary feeding programmes, 307-308
- Cottonseed cake, biological protein value, 464  
in weaning food mixtures, 394
- Creatinine, excretion in PEM, 36
- Cretinism, endemic, 83, 87, 88-89  
*see also* Goitre, endemic
- Cystine, in amino acid scoring pattern, 463
- Deaf-mutism, *see* Cretinism, endemic
- 7-Dehydrocholesterol, *see* Provitamin D
- Demineralization of bone, 469
- Dental caries, 494-496  
effect of fluoride intake on, 146, 152-154, 494  
effect of refined carbohydrates on, 303, 496  
prevalence, 495  
prevention in children, 153
- Dermatosis, criterion for classification of PEM, 532
- Diabetes, and obesity, 484
- Diarrhoea, chronic, recurrent, 206  
dehydration, 49-51  
electrolyte deficiency in, 202  
folate excretion in, 63  
underlying causes, beriberi, 139  
coeliac disease, 78  
parasitic infestations, 76  
pellagra, 131  
PEM, 29, 39, 42, 48-49, 200  
weaning, 203-207  
and vitamin A absorption, 98  
in preschool children, 205
- Dietary surveys, 260, 558  
campaign in Central America, 565-566
- Dietitian, responsibilities in nutrition education, 289-290
- 1,25-Dihydroxycholecalciferol, 112
- Diphtheria, vaccination, 269  
and vitamin deficiency, 200
- Disasters, definition, 407-408  
food shortage forecasts, 258, 259  
nutritional aspects of, 406-418  
organizational priorities, 409-411  
*see also* Famine
- DNA-synthesis, and folate deficiency, 63  
and vitamin B<sub>12</sub> deficiency, 65
- Dwarfism, nutritional, 26
- Ecology, nutritional, 162, 163-165  
nutrition and infections, 195-197
- Eggs, biological protein value, 464  
iron availability, 57, 58  
vitamin A content, 95, 107  
vitamin B content, 476  
vitamin D content, 113
- Elderly persons, energy requirements, 459
- Electrolyte metabolism, in PEM, 36, 38, 49, 202  
in severe starvation, 415
- Endemic goitre, *see* Goitre
- Energy, balance, 458  
imbalance, 486-487  
intake, in different regions, 340-342, 483  
outlook in developing countries, 344-345  
requirements, 449, 453, 457, 459  
and physical activity, 459  
and protein-energy interactions, 465-466  
resources, and food production, 14
- Enrichment, *see* Food fortification
- Environment, biological, 163
- Enzyme deficiency, congenital, in rickets, 112
- Epidemiology of nutrition, 159-192  
data collection in urban areas, 251  
incidence studies, 189  
investigative, 177-178  
descriptive, 186-188  
analytical, 188-190  
operational, 172-176  
nutrition surveys, 172, 557  
field studies, 173, 185-186

- principles, 161-169
- Ergocalciferol, 474
  - see also* Vitamin D
- Eye lesions, in riboflavine deficiency, 148
  - in vitamin A deficiency, *see* Xerophthalmia
- Faffa, 399
- Family health, 223-239
  - high risk families, 238
  - maternity care, definition, 224
  - objectives, 224-225
- Family planning, national policy, 426, 427
  - and breast feeding, 48
  - and family health, 224, 227-233
  - in health centres, 527
- Famine, 174-175
  - definition, 408
  - see also* Disasters
- Fat, intake in different countries, 483
  - malabsorption in, infections, 202-203
  - rickets, 112
  - oxidation during storage, 357-359
- Fatty acids, polyunsaturated, relation to serum lipids, 493
  - and cardiovascular disease, 489
  - and vitamin E requirement, 151
- P : S ratio, 493
- saturated, relation to serum lipids, 493
- Ferritin, 58
- Fetal life, and infections, 196
  - and maternal nutrition, 229
  - see also* Pregnancy
- FIGLU test, 63
- Field studies, 173
- Fish, iodine content, 84
  - iron availability, 58
  - liver, vitamin D content, 113
  - marketing problems, 354
  - production in developing countries, 346
  - vitamin B content, 476
- Fluorine, desirable intake, 153, 472
  - supplementation, 154
  - and dental caries, 152-154, 471
- Fluorosis, 152, 154
- Folate, 477-479
  - absorption, 64
  - body stores, 63
  - conjugated form, 478
  - metabolism, 63
  - microbiological assay, 478
  - recommended daily intake, 456
  - requirements, 478-479
  - supplementation in PEM, 41
  - and phagocyte production, 199
- Folate deficiency, 63-65, 72-74, 478
  - coeliac disease, 77
  - etiology, pathophysiology, 63-64
  - prevalence, 64-65
  - prevention, 74
  - symptoms, 68-69
  - treatment, 72
- Food, availability (supply), 340-342, 348, 422-424, 561
  - balance sheets, 340-341, 558
  - choice, and income, 244, 246, 360
  - and tradition, 225, 245
  - exports, problems in developing countries, 356-357
  - nutritional surveillance, 260
- Food additives, in weaning food mixtures, 396
- Food and nutrition policy, 218-222, 351, 419-444
  - establishment of regulations, 439
  - evaluation, 436-437
  - formulation, 432-435
  - health sector responsibilities, 437-443
  - implementation, 435-436
  - local, 441-443
  - nutrition education, 289
  - planning, 428-432
  - price control, 363-367
  - regional, 440-441
  - and biological vulnerability, 427-428
  - and food demand, 424-427
  - and food supply, 422-424
- Food fortification, 370-388
  - criteria for, 374-377
  - implementation of programmes, 385-387, 439
  - limitations, 387-388
  - nutrients, 380-382
    - ascorbic acid, 383
    - iodine, 89-91
    - iron, 74
    - niacin, 134
    - thiamine, 144
    - vitamin A, 109, 315
    - vitamin B<sub>12</sub>, 75
    - vitamin D, 115
  - philosophies, 373-374
  - target groups, 378-382
  - technological problems, 379-380, 384
  - technology, 382-385
  - vehicle foods, 383

- Food habits, 245, 284  
 significance for food donations, 412-413  
 taboos, 284-285  
 and obesity, 487  
 in rural and urban Africa, 247
- Food marketing, 353-369  
 national policy, 424  
 processed weaning foods, 401-402
- Food poisoning, 166, 175
- Food prices, indicator of nutritional status, 502  
 national policy, 425-426
- Food production, 335-352  
 effect of supplementary feeding, 309  
 energy and protein demand, 342-345  
 incentives and disincentives, 348-350  
 high-yielding crop varieties, 345-346  
 national policy, 423  
 objectives in developing countries, 345-351  
 price relationships, 349, 363  
 targets, 346-348  
 trends, 336-340  
 and population growth, 339
- Food shortage, climatic conditions, 16, 258  
 early warning systems, 258-260, 407  
 livestock production, 15  
*see also* Disasters
- Food storage, national policy, 424  
 nutritional losses during, 357-360  
 pest control, 359  
 quality control, 367-368
- Formiminoglutamic acid, excretion in folate deficiency, 63
- Fortification, *see* Food fortification
- Fruit, nutrient losses during storage, 358  
 vitamin B content, 476  
 vitamin fortification, 383
- Gamma-globulins, in PEM, 34  
 in scurvy, 124
- Genetic traits, 230-231
- German measles, and kwashiorkor, 29
- Giardia lamblia*, 76
- Glucose, infusion, malnourished children, 42  
 metabolism, and chromium, 156  
 in PEM, 36  
 rehydration solutions, 50-51
- Gluten, and coeliac disease, 77
- Goitre, 83-92  
 classification, 86  
 definition, 83  
 etiology, 84-85  
 physiopathology, 85-86  
 prevalence, 18, 86-88, 570  
 prevention, feasibility, 18  
 iodization of salt, 372  
 public health significance, 18, 88-89
- Goitrogenic compounds (goitrogens), 84
- Gomez classification, of PEM in community studies, 535
- Gossypol, 394
- Green revolution, 15
- Groundnut, vitamin B content, 476
- Groundnut cake, biological protein value, 464  
 in weaning food mixtures, 394
- Growth, and iron requirements, 59
- Growth charts, 52, 236
- Growth failure, in PEM, 44-45  
 in zinc deficiency, 471  
*see also* Weight-for-height
- Growth phases, pre- and post-natal period, 195-196
- Growth rates, relation to energy intake, 460  
 in rehabilitated children, 327
- Haematocrit, 69
- Haemoglobin, blood concentration, 56  
 estimation, 69  
 iron content, 58
- Haemolysis, red blood cells, in vitamin E deficiency, 151
- Haemopoiesis, 57
- Haemorrhages, and vitamin C deficiency, 122
- Haemosiderin, 58, 62
- Harvard standards of reference, 535
- Head circumference, relation to arm circumference, 545-546
- Health centres, mobile teams, 264, 274
- Health ministries, nutrition activities of, 438
- Health services, for family health, 234-235
- Health workers, 13  
 in family health programmes, 233-239  
 in nutrition education, 290-293, 443  
 in nutrition planning, 213-214  
*see also* Auxiliary health workers
- Heart attack, coronary, 491-492
- Heart failure, in PEM, 42, 50  
*see also* Cardiovascular disease
- Height of children effect of food deprivation in war time, 512-513  
 indicator of nutritional status, 510-514, 536-539

- see also* Weight-for-height
- Height-for-age, 538, 548-549
- Hemeralopia, *see* Night blindness
- Hepatitis, infectious, serum lipids, 202
- Hepatomegaly, criterion for classification of PEM, 532, 547
- High-risk families, 238
- Histidine, 463
- Hookworm infestations,
  - hydroxyproline excretion in, 35
  - iron deficiency in, 61, 73, 202
  - iron supplementation, therapeutic, 315 and PEM, 43
- Hydroxyproline, excretion in PEM, 35
  - index, 35
  - synthesis, effect of ascorbic acid, 200
- Hypercalcaemia, 115, 117-118, 384
- Hyperlipaemia, 485, 492
- Hyperlipoproteinaemia, 493
- Hyperparathyroidism, 113-114
- Hypertension, and cardiovascular disease, 492
  - and obesity, 485
- Hypervitaminosis A, 96, 317
  - prevention, 319, 375
  - see also* Vitamin A
- Hypoalbuminaemia, in low-paid African workers, 249
  - in PEM, 31, 34, 532
- Hypocalcaemia, in magnesium deficiency, 155, 472
  - in vitamin D deficiency, 114
- Hypoglycaemia, influence on power of concentration, 303
  - in PEM, 36
- Hypogonadism, and zinc deficiency, 154
- Hypovitaminosis A, *see* Vitamin A deficiency; Xerophthalmia
- Hypovitaminosis D, *see* Rickets; Osteomalacia
- Idiopathic steatorrhoea, *see* Coeliac disease
- Immigrants, child mortality among, 249
  - food habits, 245
  - nutritional status, 248-249
  - social situation, 244
  - in peri-urban areas, 243-255
- Immunization programmes, 216, 268-276
  - equipment, 274-275
  - mobile teams, 274
  - records, 527
  - schedules, 272-273
  - side-effects, 271
  - triple vaccine, 269
- Immunoglobulins, 271
- Inborn errors of metabolism, 230
- Incaparina, 398-400
- Income, elasticity for food, 244, 246, 360-363
  - national policy on, 425
- Industrial feeding, 303-304
- Infants, beriberi, 139-140
  - iron deficiency anaemia, 62
  - nutrient requirements, 456
    - protein, 462
    - vitamin A, 473
  - PEM, 23
  - rickets, 115-116
  - vitamin A deficiency, 108-109
  - vitamin B<sub>6</sub> deficiency, 149
  - vitamin B<sub>12</sub> deficiency, 73
  - weanling diarrhoea, 203
- Infant feeding, early bottle feeding, dangers, 30, 226
  - vegetable protein mixture (UNICEF), 40
  - see also* Baby foods; Breast feeding
- Infant formulas, for therapy of PEM, 38
- Infections, acute and chronic, 166
  - cortisone therapy, 201
  - food-borne, 166
  - herd immunity, 162, 269-270
  - immunization of children, 52, 269-272
  - mechanisms of resistance, 199-201
  - metabolic consequences, 201-202
  - mortality among children, 504
  - nitrogen loss, 202
  - prevalence in preschool children, 203-207
  - prevention, national policies, 427
  - vulnerable groups, 197
    - and family size, 231
    - and malnutrition, 28-30, 68, 193-209, 230
    - and nutrition intervention programmes, 173
    - and vitamin A deficiency, 98
- Infectious agents, and nitrogen metabolism, 201-202
  - and nutrient deficiency, 198
- Interferon, 200
- Intrinsic factor, 478
- Iodine, fortification of salt and bread, 89-90, 371-372, 572
  - iodized oil, 90
  - requirements, 471
  - thyrotoxicosis, 91
  - urinary excretion in goitre, 569-570
  - and thyroxine synthesis, 85

- Iodine deficiency, 84-92
- Iron, absorption, 58, 467  
 bioavailability, 57, 384, 466  
 distribution in the body, 58  
 food fortification with, 379-380, 384  
 programmes, 383  
 losses (in urine, faeces, menstruation), 466  
 metabolism, 57  
 adaptation to actual needs, 451  
 recommended daily intakes, 455  
 requirements, 59, 61, 451, 467  
 sources, 466  
 supplementation, 74, 315-316, 467  
 malnourished children, 40  
 toxicity, 72
- Iron deficiency, classification, 62  
 etiology, 60-62  
 prevalence, 62-63  
 prevention, 73-74  
 treatment, 71-72  
*see also* Anaemia, nutritional
- Ischaemic heart disease, *see* Cardiovascular disease
- Isoleucine, imbalance, pellagra, 126  
 in amino acid scoring pattern, 463
- Jodbasedow, 88, 91
- Keratomalacia, *see* Xerophthalmia
- Kwashiorkor, definition, 25  
 diagnostic criteria, 532  
 histopathology, 33-34  
 rehabilitation centres, 174  
 symptoms, 31, 33  
 dermatoses, 31-32, 132  
 and infections, leucocyte response, 199  
 and measles, case report, 206-208  
*see also* Protein-energy malnutrition (PEM)
- Lactation, energy requirement, 460  
 maternal malnutrition, 216  
 supplementary feeding, 300  
 nutrient requirements, 455-456  
 iron, 74  
 protein, 462  
 vitamin A, 108, 109, 474  
 promotion, 427  
*see also* Breast feeding
- Lactose intolerance, 428  
 and milk powder in weaning foods, 394  
 in PEM, 38, 39
- Land tenure systems, and food production, 350
- Leucine, imbalance, pellagra, 126  
 in amino acid scoring pattern, 463
- Linoleic acid, in weaning food mixtures, 395
- Liver, folate content, 478  
 vitamin B content, 476
- Livestock production, and affluence, 15  
 in developing countries, 346
- Lysozymes, 200
- Magnesium, 155  
 deficiency in PEM, 33, 36, 41  
 recommended daily intake, 472  
 and immunological resistance, 200
- Maize, biological protein value, 464  
 iron availability, 58  
 new varieties, 345  
 niacin enrichment, 134  
 preference to wheat, 129  
 and pellagra, 126-127, 129  
 in infant formulas (UNICEF), 40  
 in protein-rich food mixtures, 400
- Malabsorption, intestinal, 76-79  
 coeliac disease, 77-78  
 parasitic infestations, 76  
 rickets, 112  
 tropical sprue, 78-79
- Malaria, and beriberi, 202  
 and hydroxyproline excretion, 35  
 and kwashiorkor, 29, 259  
 and megaloblastic anaemia, 74
- Malnutrition, acute, 174-175  
 chronic, 165, 168  
 conditioning factors, 13-20, 163-169  
 interaction with infections, 16, 193-209  
 mechanisms, 199-203  
 maternal, and fetal death, 183, 196  
 metabolic adaptation, 451  
 prevention, 167-169  
 priorities for action, 18  
 psychological factors, 165-168  
 screening criteria for children, 416  
 statistics, 183  
 symptoms, 447  
 and child development, 17  
 and child mortality, 258  
 and school performance, 17  
 and work productivity, 19  
 and wound healing, 200  
 in industrialized countries, 194-195  
 in the weaning period, 204

- see also* Anaemia; Goitre; Protein-energy-malnutrition; Vitamin deficiencies
- Manganese, 156, 471, 472
- Marasmus, definition, 25  
 diagnostic criteria, 532  
 rehabilitation centres, 174  
 symptoms, 32  
*see also* Protein-energy malnutrition
- Marketing boards, 364
- Measles, vaccination, 269, 273  
 and kwashiorkor, 29, 202  
 and malnutrition, 206, 207-208, 270
- Meat, iodine content, 84  
 iron availability, 57  
 nutrient loss during storage, 358  
 vitamin B content, 476
- Medical training, nutrition in curriculum, 281-282
- Menarche, age of, indicator of nutritional status, 514-516
- Menstruation, iron loss, 59, 448, 467
- Mental development, and early malnutrition, 230  
 and vitamin D toxicity, 118
- Metabolic rate, basal, 448, 459  
 in undernutrition, 451
- Methionine, deficiency and collagen formation, 200  
 in amino acid scoring pattern, 463
- Milk, biological protein value, 464  
 fortification, iron, 383  
 vitamin A, 383  
 vitamin D, 115, 383  
 nutrient content, copper, 156  
 folate, 64, 74  
 iodine, 84  
 vitamin A, 95  
 vitamin D, 113  
 nutrient loss during storage, 358  
 powdered, donations, effect on breast feeding, 308  
 in weaning food mixtures, 394  
 price policy, India, 366-367  
 in PEM treatment, 38  
 in supplementary feeding, 299, 301, 308
- Millet, biological protein value, 464
- Molybdenum, 156, 471
- Morbidity, records, indicator of malnutrition, 259  
 risks, obesity, 484
- Mortality rates, analyses, nutrition surveys, 559  
 indicator of malnutrition, 502  
 and obesity, 485  
 and population structure, 165  
 and standard of living, 17, 19
- Mortality statistics, in biometric studies, 183-184
- Mucosal lesions, in coeliac disease, 78  
 in tropical enteropathy, 79
- Mycotoxins, 394  
*see also* Aflatoxin
- Myxoedema, 88
- Net protein utilization (NPU), 464
- Niacin, 475-477  
 bioavailability, 128-129, 477  
 biosynthesis from tryptophan, 475  
 equivalents, 475, 477  
 fortification, cereals, 134  
 recommended daily intake, 456, 475  
 requirements, 128
- Niacin deficiency, and antibody formation, 199  
*see also* Pellagra
- Niacytin, 128
- Nickel, 156, 171
- Nicotinic acid, *see* Niacin
- Night blindness, 97, 99, 100
- Nitrogen, catabolism in infections, 201-202  
 requirements, 461
- Nutrients, excess intake, 450
- Nutrient requirements, 445-481  
 biological variability, 448-450  
 conditioning factors, 57  
 definition, 446-448  
 metabolic adaptation, 450-452  
 recommended intakes, 450-452, 455  
 targets, 446-448
- Nutritional rehabilitation, 321-334  
 ambulatory treatment, 323-324  
 behavioural improvement, 329  
 centres, 43, 173-174, 324-333  
 day care, 325-326  
 residential, 326  
 maintenance cost, 329-330  
 organization, 331  
 hospital treatment, 322-323  
 limitations, 332  
 weight gain, as indicator, 327-328  
 in times of disaster, 413, 414-417
- Nutritional status, appraisal of, 556-575  
 criteria, 530-555  
 improvement, feeding programmes, 217, 305

- immunization, 270  
 indicators, 500-519, 205  
 in urban populations, 247-250  
 of children, 248  
*see also* Anthropometry; Biometry; Surveys, nutritional
- Nutrition clinics, 323-324
- Nutrition education, approaches, 282-285  
 coordination of activities, 279, 287, 288, 292  
 interagency collaboration, 236  
 national policies, 426, 439, 443  
 objectives, 277-280  
 prevention of nutritional anaemias, 73  
 programme planning, 285-287  
 research and evaluation, 293-294  
 responsibility of the dietitian, 289-290  
 techniques, 287  
 target groups, 280-282  
 for persons requiring therapeutic diets, 280  
 for schoolchildren, 237, 302  
 in rehabilitation centres, 325-330
- Nutrition intervention programmes, assessment of nutritional status, 219  
 coverage, 220-221  
 evaluation, 177  
 pilot projects, 219-220  
 planning and organization, 213-222  
 timing, 219  
 for peri-urban areas, 250-255  
*see also* Food fortification; Supplementary feeding
- Nyctalopia, *see* Night blindness
- Oatmeal, vitamin B content, 476
- Obesity, 340, 483-489  
 etiology, 485, 488  
 health risks, 485-486  
 and carbohydrate metabolism, 484  
 and cardiovascular disease, 492  
 and physical activity, 487  
 in children, 488
- Oedema, criterion for classification of PEM, 531-532  
 and kwashiorkor, 25, 31  
 and sodium metabolism, 36  
 and starvation, 414, 416  
 and susceptibility to infections, 199
- Oilseed protein, in weaning food mixtures, 392, 393-394
- Osteomalacia, 111-119, 474  
*see also* Rickets
- Osteoporosis, 469
- Overweight, *see* Obesity
- Pan American Health Organization (PAHO), 106
- Pantothenic acid deficiency, antibody formation in, 199, 200
- Parasitic infestations, and anaemia, 61-62, 67, 76  
 and pellagra, 130
- Peas, vitamin B content, 476
- Pellagra, 13, 126-135  
 clinical signs, 130-131  
 control and prevention, 133-134  
 diet therapy, 133  
 niacin administration, 133  
 differential diagnosis, 131-133  
 epidemiology, 127-130  
 mental symptoms, 131  
 niacin availability, 477  
 in urban populations, 250
- Perinatal death, 183  
*see also* Malnutrition, maternal
- Peri-urban areas, 243-255  
 actions to improve the nutrition situation, 253-255  
 collection of statistical data, 250-253  
 factors influencing food choice, 243-247  
 hygienic conditions, 246  
 nutritional status, adults, 248-249  
 children, 248
- Pertussis, *see* Whooping-cough
- Pest control, in food storage, 359
- Phenylalanine, in amino acid scoring pattern, 463
- Phosphatase, alkaline, and vitamin D deficiency, 114
- Phosphate, and iron absorption, 58
- Phosphorus requirements, 470
- Physical activity, adaptation to undernutrition, 19  
 influence of anaemia, 68
- Phytic acid, and calcium absorption, 113  
 and iron absorption, 58  
 and zinc absorption, 471
- Pilot projects, for nutrition programmes, 220
- Poliomyelitis, vaccination, 269, 273
- Polyneuritis, and thiamine deficiency, 138
- Population, definition in epidemiology, 163, 180  
 growth, and food demand, 339, 343  
 national policy 426

- Pork, vitamin B content, 476
- Post-mortem examinations, for nutrition surveys, 559
- Potassium, deficiency in diarrhoea, 49, 50  
metabolism in PEM, 36  
supplementation for infants, 39  
in rehydration solutions, 50-51
- Pregnancy, hemeralopia, 97  
infections, 196, 230  
intervention programmes, 216-217  
  folate supplementation, 75, 479  
  iron supplementation, 74, 317, 468  
  supplementary feeding, 46, 300  
malnutrition, folate deficiency, 65, 479  
  iron deficiency, 62, 379, 467  
  PEM, 196  
  vitamin B<sub>6</sub> deficiency, 150  
nutritional demands, 227-229, 455-456  
  energy, 460  
  iron, 59-60, 467  
  protein, 462  
spacing, 229  
  national policy, 427  
teenage, 196, 230  
vitamin D toxicity, 118
- Preschool children, diarrhoea, 203-207  
malnutrition, intervention programmes, 215-222  
  supplementary feeding, 298-299, 306  
in family health activities, 237  
*see also* Weaning
- Prices policy, 363-367  
  government intervention on retail prices, 355, 363  
  influence on agricultural production, 349, 354  
  subsidies, 365-366
- Pronutro, 400
- Properdin, 200
- Protein, biological value, 463-465  
  efficiency, 461  
  food fortification, 383, 385  
  intake, average, 454  
  recommended, 455  
  metabolism, infections, 202  
  interaction with energy, 465-466  
  requirements, 454, 460-462  
  sources, unconventional, 391, 394
- Protein-Calorie Advisory Group (PAG), 184, 221
- Protein-energy malnutrition (PEM), 23-54, 465  
  classification, 26-27, 530-555  
  grading severity, 549-550  
  control, 45-52, 167-169  
  food fortification, 381  
  intervention programmes, 215  
  rehabilitation centres, 174, 324-333  
  recipes, 38-41  
  tube feeding, 40  
epidemic, 175  
epidemiology, 15, 28-31, 172-174  
hypothermia, 41-42  
indicators, diagnosis, 37, 501-529, 532  
  body weight, 528  
mortality rates, 322  
physiopathology, 33-37  
  carbohydrate metabolism, 36  
  electrolytes, 36  
  pancreatic secretion, 37  
prevalence, 27-28, 45, 298-299  
relapses, 322  
symptoms, 31-33  
  subclinical, 448  
social significance, 18  
terminology, 24-26  
  and antibody formation, 199  
  and artificial infant feeding, 30  
  and immunization, 270-272  
  and infections, 29, 195-209, 259  
  and magnesium deficiency, 155  
  and parasitic infestations, 43  
  and pregnancy, 196  
  and vitamin A deficiency, 98  
  in Central America, survey of, 569  
  in urban areas, 248
- Protein-rich foods, 308  
  expenditure on, in different regions, 362  
  financial subsidies for, 365
- Protein supply, outlook in developing countries, 344  
  tables for different regions, 340-341
- Prothrombin, 480
- Provitamin A, *see*  $\beta$ -Carotene
- Provitamin D, 112
- Pteroylmonoglutamic acid, *see* Folate
- Pyridoxine, *see* Vitamin B<sub>6</sub>
- QUAC stick, 261, 545
- Reference population, for nutritional standards, 458-459, 535
- Rehydration, centres, 51  
  in diarrhoea, 50-51
- Renal insufficiency, and obesity, 485  
  and rickets, 112

- Respiratory diseases, mortality in children, 504
- Retinol, *see* Vitamin A
- Riboflavin, 475-477  
recommended daily intake, 456, 475  
requirements, 148
- Riboflavin deficiency, 146-149, 477  
antibody formation, 199  
epidemiology, 147  
symptoms, 147-148  
therapy, 148  
in PEM, 33
- Rice, biological protein value, 464  
fortification with vitamins, technology, 382  
iron availability, 58  
polished thiamine content, 136, 140, 177  
prices policy, different countries, 366  
protein-rich, 346  
vitamin B content, 476
- Rickets, 111-119, 474  
clinical signs, 114-115  
history, 111-112  
hypocalcaemia, 114  
physiopathology, 112-113  
prevalence, 113  
prevention, 115-118  
vitamin D supplementation, 315  
and calcium intake, 113  
and infections, 198  
in peri-urban areas, 248  
*see also* Osteomalacia
- Rubella, in pregnancy, 196
- Rye, vitamin B content, 476
- Salt, iodized, 89-90, 372
- Sample selection, for nutritional surveys, 561
- Schistosomiasis, and pellagra, 130
- School feeding, 301-303, 306, 442
- School health, 237
- Scurvy, 13, 120-125, 479  
antibody formation, 124  
biochemical signs, 123-124  
clinical signs, 122-123, 200, 479  
history, epidemiological, 120-121  
infections, 198  
prevention, 124-125
- Selenium, 156, 471
- Serum lipids, 485, 489-490  
and diet composition, 493
- Sesame, biological protein value, 464  
vitamin B content, 476
- Shanty towns, *see* Peri-urban areas
- Silicon, 471
- Single cell protein, 394
- Sitosterol, and serum lipids, 493
- Skeleton, calcification, defective, in vitamin C deficiency, 123  
deminceralization in vitamin D deficiency, 114, 116  
fluorosis, 152  
maturation as indicator of nutritional status, 538-539
- Skinfold thickness, in obesity, 484
- Slums, *see* Peri-urban areas
- Smallpox, vaccination, 269, 273
- Sodium, loss in diarrhoea, 49  
metabolism in PEM, 36  
in rehydration solutions, 50-51
- Sorghum, amino acid imbalance, pellagra, 127
- Soya bean (soybean), biological protein value, 464  
goitrogens, 84  
iron availability, 58, 467  
vitamin B content, 476  
in infant formulas, 40  
in weaning food mixtures, 391, 394
- Spinach, iodine source, 84  
iron availability, 58  
vitamin B content, 476
- Splenomegaly, criterion for classification of PEM, 547-548
- Staple foods, 16
- Starch, intake, different countries, 483
- Starvation, refeeding, 414-417  
*see also* Disaster; Famine
- Steatorrhoea, and coeliac disease, 77
- Strongyloides stercoralis*, 76
- Stunting, 537-538
- Subsidy systems, 365-367
- Sucrose, fortification with vitamin A, 109, 385  
intake, in different countries, 483  
and cardiovascular disease, 490  
and dental caries, 494-496  
and vitamin B deficiency, 477  
intolerance in infants, special formulas, 39
- Sugar, *see* Sucrose
- Superamine, 399
- Supplementary feeding, 297-312, 324  
benefits and limitations, 304-309  
children, after immunization, 271  
cost-effectiveness, 307-308

- coverage, 306
- foods, 300-301
- historical review, 297-298
- influence on food production, 309
- national policy, 426, 442
- nutrition education, 288, 308-309
- objectives, 298
- organization, 299
- price policy, 365
- schools, 301-303
- Supplementation programmes, vitamins, minerals, 313-320
- Surveillance, nutritional, 176-177
  - activities, local level, 442
  - coverage, 263-264
  - methods, 258-263
    - weight charts, 520-529
  - principles, 256-257
  - reporting, 265-266
  - schoolchildren, 529
  - staff, 264-265
  - and nutrition education, 288
  - in national food and nutrition policies, 437, 438
  - in times of disaster, 413-414
- Surveys, nutritional, 186-187, 261-263, 556-575
  - criteria, 219
  - evaluation, 173, 562-563
  - example survey in Central America, 563-574
  - methodology, 558-560
    - biochemical tests, 261-262
    - clinical examination, 261
    - blood tests, 70
    - QUAC stick method, 261
  - objectives, 556-558
  - reporting, 563
  - sampling, 560-562
    - group selection, 172
  - see also* Dietary surveys; Epidemiology
- Systems analysis, and epidemiological interpretation, 190-191
  - in nutrition intervention programmes, 221
- Tapeworm infestation, vitamin B<sub>12</sub> deficiency, 67
- Tetanus, vaccination, 269, 273
- Therapeutic diets, 280
- Thiaminase, 144
- Thiamine, food fortification with, 383
  - losses during food storage, cereals, 358
  - meat, 358
  - recommended daily intake, 456, 475
- Thiamine deficiency, and antibody formation, 199
  - see also* Beriberi
- Threonine, in amino acid scoring pattern, 463
- Thyroid enlargement, *see* Goitre, endemic
- Thyrotoxicosis, *see* Jodbasedow
- Thyroxine, and goitre, 85
- Tin, 156, 471
- Tissue, adipose, cell number and obesity, 486, 488
- Tocopherol, *see* Vitamin E
- Tortillas, availability of niacin, 129
- Trace elements, 155-157, 470-472
  - relation to cardiovascular disease, 490
  - toxicity levels, 450
- Transferrin, 35, 58, 62
  - degree of saturation, iron status, 71
- Transketolase activity, and thiamine deficiency, 139
- Trichuris trichiura*, infestation, and iron deficiency, 61, 73
- Triglycerides, serum, and carbohydrate intake, 493
  - and cardiovascular disease, 490
- Tropical enteropathy, 79
- Tropical sprue, 78-79
  - and iron absorption, 60
- Trypsin inhibitor, in soya beans, 394
- Tryptophan, conversion to niacin, 126, 475
- deficiency, and antibody formation, 199
  - in amino acid scoring pattern, 463
- Tube feeding, of marasmic children, 40
- Tuberculosis, vaccination, 269, 273
  - and kwashiorkor, 29, 270
  - tuberculin reactions, 199
  - in African miners, 249
- Tyrosine, in amino acid scoring pattern, 463
- Undernutrition, *see* Malnutrition
- Unsaturated iron binding capacity (UIBC), 62
- Urban areas, 240-255
  - factors influencing food choice in, 243-247
  - food habits, Africa, 247
  - food retail systems, 355
  - growth rates, 242, 354

- percentage of country's inhabitants living in, 241
- sex disparity in, 246
- Vaccination schedules, 272-273
  - for children, 269, 273
  - see also* Immunization programmes
- Valine, in amino acid scoring pattern, 463
- Vanadium, 156, 471
- Veal, iron availability, 58
- Vegetables, vitamin B content, 476
- Vegetarians, vitamin B<sub>12</sub> deficiency in, 75
- Vitamin deficiencies, antibody formation in, 199, 200
  - biochemical tests, 262
  - in urban populations, 250
  - see also* Beriberi; Pellagra; Rickets; Scurvy; Xerophthalmia
- Vitamin supplementation, 313-320
- Vitamin A, food fortification, technological problems, 383
  - injection, 106
  - intakes, 95-96, 472
  - losses during food storage, 357
  - recommended intake, 108, 456
  - requirements, 472-473
  - retinol equivalents, 473
  - sources in the diet, 95, 107
  - supplementation, oral, 106, 317
- Vitamin A deficiency, 94-110
  - eye infections, 105
  - infectious disease, 98, 198
  - and antibody formation, 199, 200
  - and PEM, 33
  - and socioeconomic conditions, 96, 107
  - in childhood, 96-97
  - in pregnancy, 97
  - see also* Xerophthalmia
- Vitamin B<sub>1</sub>, *see* Thiamine
- Vitamin B<sub>2</sub>, *see* Riboflavin
- Vitamin B<sub>6</sub>, antagonists, 149-150
  - indicator of nutritional status, 150
  - physiology, 149
  - requirements, 149, 480
- Vitamin B<sub>6</sub> deficiency, 149-150
  - and oral contraceptives, 150
- Vitamin B<sub>12</sub>, 477-479
  - absorption, 67
  - metabolism, 65-66
  - recommended intake, 456
  - requirements, 66
- Vitamin B<sub>12</sub> deficiency, 64-65, 67-69
  - symptoms, 68-69
  - treatment, supplementation, 72-73, 75
- Vitamin C, *see* Ascorbic acid
- Vitamin D, 474-475
  - content of foods, 113
  - food fortification with, 383-384
  - metabolism, 112-113
    - interaction with calcium, 113, 474
  - recommended daily intake, 456
  - supplementation, oral, 317
  - synthesis in the skin, 113
  - toxicity, 117-118
- Vitamin D deficiency, *see* Osteomalacia; Rickets
- Vitamin E, 151-152, 480
  - requirements, influence of polyunsaturated fatty acids on, 151
- Vitamin E deficiency, haemolysis test for, 151
- Vitamin K, 480
- Vitasoy, 400
- Vulnerability, biological, 427
- Water, 457
  - fluoridation, 153
  - metabolism in PEM, 36
  - supply in times of disaster, 411
- Weaning, 30, 390
  - and PEM, death rate, 506
  - diarrhoea, 203-207
  - practices, 204
- Weaning food mixtures, 389-405
  - composition, 393-396
  - fortification, 379-380
  - local mixes, 48
  - promotion, 400
  - protein-rich, 48, 291-292
  - retail price, 401
  - testing, 397-398
  - types and characteristics, 391-393
  - see also* Baby foods
- Weight charts, 520-529
  - design, 521-522
- Weight-for-age, 535-536, 549
  - indicator of nutritional status, 219, 262
  - influence of malnutrition, diarrhoea and infections, 205, 206
  - PEM, 44-45
  - recovery in rehabilitation centres, 327
  - see also* Anthropometry
- Weight-for-height, 540-545, 549
  - age dependency, 539
  - indicator of nutritional status, 262, 414, 501, 537

- mathematical relationship, 543-545
  - see also* Anthropometry
- Wellcome classification, for PEM, 532-533
- Wheat, biological protein value, 464
  - coeliac disease, 77
  - iron, availability, 58
  - fortification with, 74
  - vitamin B content, 476
  - in supplementary feeding programmes, 308
- Wheat-soya-blend (WSB), 399
- Whooping-cough, vaccination, 269, 273
  - and kwashiorkor, 29, 202, 270
- World Food Programme (WFP), 304, 309, 324
- Wound healing, 200
  - and ascorbic acid, 479
- Xerophthalmia, 94-110
  - clinical features, 99-105
  - blindness, 97
- epidemiology, 96-98
  - climatic conditions, 97
  - physiopathology, 94-96, 98
  - prevention, 18, 107-109
    - milk fortification, 315
  - sex differences, 97
  - social significance, 18, 98
  - survival rate, 97
  - treatment, 105-107
    - rehabilitation centres, 326-327
    - and breast feeding, 474
  - and PEM, 40, 95, 98, 202
  - see also* Vitamin A deficiency
- Xerosis, 100-104
  - skin, chronic malnutrition, 132
- Xylose-absorption test, and tropical enteropathy, 79
- Zinc, 154-155, 471
  - antagonism to copper, 156
  - deficiency, 155
  - requirements, 155